

Walmart Inc.

-PE Product Labels –

n



Let and Expiration Mo.
No Prim/No Varnish

PARENTS:
www.hugobosch.com
TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

Drug Facts (continued)

Warnings: This product contains acetaminophen. Severe liver damage may occur if you take more than 4,000 mg of acetaminophen in 24 hours.

Do not use: with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if you have:

- liver disease
- kidney disease
- heart disease
- thyroid disease
- difficulty in urination due to enlargement of the prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- high blood pressure

Drug Facts (continued)

Directions:

- adults and children 12 years and over: take 2 caplets every 4 hours
- children under 12 years: ask a doctor

Other information:

- each caplet contains: sodium 3 mg
- TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN

Questions or comments? 1-888-287-1915

cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, sucralose, talc, titanium dioxide

Satisfaction guaranteed - For questions or comments please call 1-888-287-1915

DISTRIBUTED BY: Walmart Inc., Bentonville, AR 72716
This product is not manufactured or distributed by Johnson & Johnson Corporation, owner of the registered trademark Tylenol® Cold + Flu Severe.

729665
6 81131 18718 3

equate™

Compare to Tylenol® Cold + Flu Severe active ingredients*

SEVERE

Cold & Flu

Acetaminophen - Pain Reliever/ Fever Reducer

Dextromethorphan HBr - Cough Suppressant

Guaifenesin - Expectorant

Phenylephrine HCl - Nasal Decongestant

Relieves:

- Headache & body aches
- Fever & sore throat
- Cough
- Nasal congestion
- Mucus & chest congestion

24 CAPLETS

Actual Size

Drug Facts

Active ingredients (in each caplet)

Purpose

Acetaminophen 325 mg Pain reliever/fever reducer
Dextromethorphan HBr 10 mg Cough suppressant
Guaifenesin 200 mg Expectorant
Phenylephrine HCl 5 mg Nasal decongestant

Drug Facts (continued)

Uses: temporarily relieves these common cold and flu symptoms:

- headache
- cough
- nasal congestion
- sore throat
- minor aches and pains
- helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive
- temporarily reduces fever

Drug Facts (continued)

When using this product do not exceed recommended dosage.


Stop use and ask a doctor if:

- new symptoms occur
- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any symptoms.

B-2203-503A-08-FLU
REV0922C50308


Equate 44-503A



[illegible]

Day & Night
50844 REV0517A61569409
Bentleyville, AR 72716
DISTRIBUTED BY: Walmart Inc.,
Bentleyville, AR 72716
This product is not manufactured or distributed by
Roche/Boehringer LLC, owner of the registered
trademark Maximum Strength SINUS-MAX®

Satisfaction guaranteed - Or we'll
replace it or give you your money
back. For questions or comments or
to report an undesired reaction or
side effect, please call
1-888-287-1915.



PEEL HERE FOR MORE DRUG FACTS

how2recycle.info
PAPER TRAY
MULTI-LAYER
CARD

Do not take the **Daytime** and **Nighttime** caplets at the same time.

Drug Facts (continued)

Warnings: This product contains acetaminophen. Severe liver damage may occur if you take more than 4,000 mg of acetaminophen in 24 hours.

Do not use:

- If you have ever had an allergic reaction to this product or any of its ingredients
- If you have ever had an allergic reaction to this product or any of its ingredients

Other information:

- If you are taking other drugs containing acetaminophen, ask a doctor or pharmacist before taking this product.
- If you are taking other drugs containing acetaminophen, ask a doctor or pharmacist before taking this product.

Drug Facts

Compare to Maximum Strength Mucinex® SINUS-MAX® Day & Night Active Ingredients*

Daytime (Sinus Day)	Nighttime (Sinus Night)
Guafenesin 200 mg	Phenylephrine HCl 5 mg
Acetaminophen 325 mg	Phenylephrine HCl 5 mg
Pain reliever	Nasal decongestant
Purpose	Purpose
Relieves sinus pressure, headache & congestion	Relieves sinus pressure, headache & congestion
Thins & loosens mucus	Thins & loosens mucus

Do not use:

- If you are taking other drugs containing acetaminophen, ask a doctor or pharmacist before taking this product.
- If you are taking other drugs containing acetaminophen, ask a doctor or pharmacist before taking this product.

Other information:

- If you are taking other drugs containing acetaminophen, ask a doctor or pharmacist before taking this product.
- If you are taking other drugs containing acetaminophen, ask a doctor or pharmacist before taking this product.

equate™

Maximum Strength
Day & Night Sinus

equate™

Maximum Strength
Day & Night Sinus

Compare to Maximum Strength Mucinex® SINUS-MAX® Day & Night Active Ingredients*

Day

Acetaminophen - Pain Reliever,
Guafenesin - Expectorant,
Phenylephrine HCl - Nasal Decongestant

- Relieves sinus pressure, headache & congestion
- Thins & loosens mucus

10 DAYTIME CAPLETS

Night

Acetaminophen - Pain Reliever,
Diphenhydramine HCl - Antihistamine/
Cough suppressant,
Phenylephrine HCl - Nasal Decongestant

- Relieves nasal congestion, sinus pressure & pain
- Runny nose, sneezing & cough

10 NIGHTTIME CAPLETS

Actual Size

Drug Facts (continued)

Ask a doctor before use if you have:

- heart disease
- liver disease
- difficulty in urination due to enlargement of the prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- a cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- glaucoma (Nighttime only)

Ask a doctor or pharmacist before use if you are:

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers (Nighttime only)

When using this product:

- do not exceed recommended dosage
- excitability may occur, especially in children (Nighttime only)
- marked drowsiness may occur (Nighttime only)
- alcohol, sedatives, and tranquilizers may increase drowsiness (Nighttime only)

Drug Facts (continued)

Stop use and ask a doctor if:

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

If taking NIGHTTIME and DAYTIME products, carefully read each section to ensure correct dosing. Do not take DAY & NIGHT at the same time.

Drug Facts (continued)

Directions

- do not use more than directed
- do not take more than 12 caplets in any 24-hour period
- adults and children 12 years and over: take 2 caplets every 4 hours
- children under 12 years: do not use

Other information

- TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

Inactive ingredients (Daytime only)

corn starch, croscarmellose sodium, croscarmellose, FD&C blue #1 aluminum lake, FD&C blue #2 aluminum lake, iron oxide yellow, magnesium stearate, methacrylic acid and ethyl acrylate copolymer, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, sodium bicarbonate, stearic acid, talc, titanium dioxide

Drug Facts (continued)

Inactive ingredients (Nighttime only)

corn starch, croscarmellose sodium, croscarmellose, FD&C blue #1 aluminum lake, FD&C blue #2 aluminum lake, iron oxide yellow, magnesium stearate, methacrylic acid and ethyl acrylate copolymer, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, sodium bicarbonate, stearic acid, talc, titanium dioxide

Questions or comments? 1-888-287-1915

50844
REV0517A61569409
R-2203-615694-09SM-DF

DAY & NIGHT TWIN PACK



MAXIMUM STRENGTH
Sinus Relief
 Acetaminophen 325mg
 Dextromethorphan HBr 1mg
 Guafenesin 500 mg
 Pseudoephedrine HCl 5mg

MAXIMUM STRENGTH
Sinus Relief
 Acetaminophen 325mg
 Dextromethorphan HBr 1mg
 Doxylamine succinate 4.2mg
 Pseudoephedrine HCl 5mg

DAY & NIGHT TWIN PACK


NDC 055-154-01

Compare
to Mucinex®
Sinus-Max® Day
and Night
Maximum
Strength Active
Ingredients*

Maximum Strength**

Fast Dissolving Softgels
Sinus Relief

Day



ACETAMINOPHEN
Pain Reliever

Dextromethorphan HBr
Cough Suppressant

Guafenesin
Expectorant

Phenylephrine HCl
Nasal Decongestant

- Relieves sinus pressure, headache & congestion
- Controls cough
- Thins & loosens mucus



Adults

16 SOFTGELS
(Liquid Filled Capsules)

Night



ACETAMINOPHEN
Pain Reliever

Dextromethorphan HBr
Cough Suppressant

Doxylamine succinate
Antihistamine

Phenylephrine HCl
Nasal Decongestant

- Relieves nasal congestion, sinus pressure & pain
- Controls cough
- Relieves runny nose & sneezing



Adults

8 SOFTGELS
(Liquid Filled Capsules)

24 TOTAL
AGES 12+

LOT NO:

EXP DATE:

COATING
FREE AREA



6 81131 27526 2

PARENTS:
Learn about your child's illness
www.StopMedicineAbuse.org



COATED PAPER MULTI-LAYER BOX TRAY



Tamper evident: Do not use if carton is damaged or if printed text on blister is broken or missing.

Drug Facts

Active ingredients (in each Softgel) Purposes

Sinus Daytime

Acetaminophen 325 mg.....Pain reliever

Dextromethorphan HBr 1 mg.....Cough suppressant

Guafenesin 500 mg.....Expectorant

Phenylephrine HCl 5 mg.....Nasal decongestant

Sinus Nighttime

Acetaminophen 325 mg.....Pain reliever

Dextromethorphan HBr 1 mg.....Cough suppressant

Doxylamine succinate 4.2 mg.....Antihistamine

Phenylephrine HCl 5 mg.....Nasal decongestant

Uses ■ temporarily relieves:

- nasal congestion
- headache
- cough
- minor aches and pains
- sinus congestion and pressure
- runny nose and sneezing (Nighttime only)
- temporarily promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to relieve the bronchial passageways of bothersome mucus and make coughs more productive (Daytime only)

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 12 softgels in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

■ If any drowsiness occurs, stop use and seek medical help right away.

■ If a skin reaction occurs, stop use and seek medical help right away.

Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

■ If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- diabetes
- high blood pressure
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- glaucoma (Nighttime only)
- breathing problems such as emphysema or chronic bronchitis (Nighttime only)

Drug Facts (continued)

■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

■ cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers (Nighttime only)

When using this product

- do not use more than directed
- excitability may occur, especially in children (Nighttime only)
- marked drowsiness may occur (Nighttime only)
- alcohol, sedatives, and tranquilizers may increase drowsiness (Nighttime only)
- avoid alcoholic drinks (Nighttime only)
- be careful when driving a motor vehicle or operating machinery (Nighttime only)

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 12 softgels in any 24-hour period
- adults and children 12 years of age and older: take 2 softgels every 4 hours
- children under 12 years of age: do not use

Other Information

- store at room temperature 15°-30°C (59°-86°F) and avoid excessive heat.

Inactive ingredients (Daytime only) FD&C yellow #6, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, partial carbon solution, and white ink.

(Nighttime only) FD&C yellow #10, FD&C blue #1, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, partial carbon solution, and white ink.

Questions or comments? 1-800-547-0995

DAY & NIGHT TWIN PACK



MAXIMUM STRENGTH
Daytime
Severe Cold & Flu
ACETAMINOPHEN 325mg
Dextromethorphen HBr 10mg
Guaifenesin 200mg
Phenylephrine HCl 5mg

MAXIMUM STRENGTH
Nighttime
Severe Cold & Flu
ACETAMINOPHEN 325mg
Dextromethorphen HBr 10mg
Doxylamine succinate 6.25mg
Phenylephrine HCl 5mg

DAY & NIGHT TWIN PACK



NDC 48035-586-04

Compare to Mucinex® Fast-Max® Day Severe Cold and Night Cold & Flu Active Ingredients*

Maximum Strength**

Fast Dissolving Softgels

Severe Cold & Flu

Day Severe Cold

ACETAMINOPHEN
Pain Reliever / Fever Reducer



Dextromethorphan HBr
Cough Suppressant

Guaifenesin
Expectorant

Phenylephrine HCl
Nasal Decongestant

- Relieves aches, fever & sore throat
- Controls cough
- Relieves nasal & chest congestion
- Thins & loosens mucus



Actual Size

16 SOFTGELS
(Liquid Filled Capsules)

24 TOTAL
AGES 12+

Night Cold & Flu

ACETAMINOPHEN
Pain Reliever / Fever Reducer



Dextromethorphan HBr
Cough Suppressant

Doxylamine succinate
Antihistamine

Phenylephrine HCl
Nasal Decongestant

- Relieves aches, fever & sore throat
- Controls cough
- Relieves nasal congestion
- Relieves runny nose & sneezing



Actual Size

8 SOFTGELS
(Liquid Filled Capsules)

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org



how2recycle.info

COATED PAPER BOX

MULTI-LAYER TRAY

*Not recycled in all communities

****Per 4-hour dose**

Do not take the Day Severe Cold and Night Cold & Flu softgels at the same time. Do not take more than a total of 12 softgels in a 24-hour period.

Do not take the first dose of the Night Cold & Flu softgels sooner than 4 hours after the last dose of the Day Severe Cold softgels unless directed by a doctor.

Take only as directed.

Keep carton for complete product information

tamper evident: Do not use if carton is damaged or if printed seal on blister is broken or missing.

DISTRIBUTED BY: Walmart Inc.,
Bentonville, AR 72716
PRODUCT OF CHINA

*This product is not manufactured or distributed
by Reckitt Benckiser LLC, owner of the
registered trademarks Mucinex® Fast-Max®
Day Severe Cold and Night Cold & Flu.

Satisfaction guaranteed - For questions
or comments please call 1-888-287-1915.

WM23-00

Drug Facts

Active ingredients (in each Softgel) Purposes

Daytime Severe Cold
Acetaminophen 325 mg.....Pain reliever/fever reducer
Dextromethorphan HBr 10 mg.....Cough suppressant
Guaifenesin 200 mg.....Expectorant
Phenylephrine HCl 5 mg.....Nasal decongestant

Active ingredients (in each Softgel) Purposes

Nighttime Cold & Flu
Acetaminophen 325 mg.....Pain reliever/fever reducer
Dextromethorphan HBr 10 mg.....Cough suppressant
Doxylamine succinate 6.25 mg.....Antihistamine
Phenylephrine HCl 5 mg.....Nasal decongestant

Uses

- temporarily relieves these common cold and flu symptoms:
 - cough
 - minor aches and pains
 - headache
 - nasal congestion
 - sore throat
 - runny nose and sneezing (Nighttime only)
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive. (Daytime only)
- controls cough to help you get to sleep (Nighttime only)
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 12 softgels in 24 hours, which is the maximum daily amount

- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- diabetes
- high blood pressure
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- glaucoma (Nighttime only)

Drug Facts (continued)

- breathing problems such as emphysema or chronic bronchitis (Nighttime only)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers (Nighttime only)

When using this product

- do not use more than directed
- excitability may occur, especially in children (Nighttime only)
- marked drowsiness may occur (Nighttime only)
- alcohol, sedatives, and tranquilizers may increase drowsiness (Nighttime only)
- avoid alcoholic drinks (Nighttime only)
- be careful when driving a motor vehicle or operating machinery (Nighttime only)

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away.

Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions


- do not take more than directed (see Overdose warning)
- do not take more than 12 softgels in any 24-hour period
- adults and children 12 years of age and older: take 2 softgels every 4 hours
- children under 12 years of age: do not use

Other information store at room temperature 15°-30°C (59°-86°F) and avoid excessive heat.

Inactive ingredients (Daytime only) FD&C yellow #6, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol sorbitan solution, and white ink
(Nighttime only) D&C yellow #10, FD&C blue #1, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol sorbitan solution, and white ink

Questions or comments? 1-888-287-1915





NDC 48035-082-34

equate™

**children's
COLD &
COUGH**

Brompheniramine Maleate
Antihistamine
Dextromethorphan HBr
Cough Suppressant
Phenylephrine HCl
Nasal Decongestant

6 YEARS AND OLDER
Relief of:

- ☐ Nasal congestion
- ☐ Runny nose
- ☐ Itchy, watery eyes
- ☐ Coughing, sneezing

■ Alcohol Free

**Grape
Flavored**

8 FL OZ (237 mL)



Dosage Cup
included



NDC 48035-082-34

equate™

**children's
COLD &
COUGH**

Brompheniramine Maleate
Antihistamine
Dextromethorphan HBr
Cough Suppressant
Phenylephrine HCl
Nasal Decongestant

6 YEARS AND OLDER
Relief of:

- ☐ Nasal congestion
- ☐ Runny nose
- ☐ Itchy, watery eyes
- ☐ Coughing, sneezing

■ Alcohol Free

**Grape
Flavored**

8 FL OZ (237 mL)



Dosage Cup
included

Grape Flavored

6 YEARS AND OLDER

COLD & COUGH


equate™
Drug Facts

Active ingredients (in each 10 mL) **Purposes**
Brompheniramine maleate, USP 2 mg.....Antihistamine
Dextromethorphan HBr, USP 10 mg.....Cough suppressant
Phenylephrine HCl, USP 5 mg.....Nasal decongestant

Uses

- temporarily relieves cough due to minor throat and bronchial irritation occurring with a cold, and nasal congestion due to the common cold, hay fever or other upper respiratory allergies
- temporarily relieves these symptoms due to hay fever (allergic rhinitis):
 - runny nose
 - sneezing
 - itchy, watery eyes
 - itching of the nose or throat
- temporarily restores freer breathing through the nose

Warnings

- Do not use
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI), certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease, or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- to make a child sleepy

Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- cough that lasts or is chronic such as occurs with smoking, asthma or emphysema

Ask a doctor or pharmacist before use if you are taking

- sedatives or tranquilizers
- When using this product
 - do not use more than directed
 - may cause marked drowsiness
 - avoid alcoholic beverages
 - alcohol, sedatives, and tranquilizers may increase drowsiness
 - be careful when driving a motor vehicle or operating machinery
 - excitability may occur, especially in children

Stop use and ask a doctor if

- you get nervous, dizzy, or sleepless
- symptoms do not get better within 7 days or are accompanied by fever
- cough lasts for more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. A persistent cough may be a sign of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Drug Facts (continued)
Directions

- do not take more than 6 doses in any 24-hour period
- measure only with dosage cup provided
- keep dosage cup with product
- mL = milliliter

age	dose
adults and children 12 years and over	20 mL every 4 hours
children 6 to under 12 years	10 mL every 4 hours
children under 6 years	do not use

Other information

- each 10 mL contains: sodium 4 mg
- store at 20°-25°C (68°-77°F)

Inactive ingredients anhydrous citric acid, edetate disodium, FD&C blue no. 1, FD&C red no. 40, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sorbitol solution

Questions or comments? 1-888-287-1915

DISTRIBUTED BY: Wal-Mart Stores, Inc.,

Bentonville, AR 72716

*This product is not manufactured or distributed by Wyeth LLC, owner of the registered trademark Dimetapp®.

**DO NOT USE IF PRINTED NECKBAND IS
BROKEN OR MISSING**



Satisfaction guaranteed -
Or we'll replace it or give you
your money back. For questions
or comments or to report an
undesired reaction or side effect,
please call 1-888-287-1915.

Gluten-Free

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org




how2recycle.info



LOT NO.

EXP.

198734 2C C9

**Drug Facts**

Active Ingredients (in each 5 mL) **Purposes**
 Dextromethorphan HBr 5 mg.....Cough suppressant
 Guaifenesin 100 mg.....Expectorant
 Phenylephrine HCl 2.5 mg.....Nasal decongestant

Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily relieves:
 - cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
 - the intensity of coughing
 - the impulse to cough to help your child get to sleep
 - nasal congestion due to a cold
 - stuffy nose

Warnings

Do not use in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.

Ask a doctor before use if the child has

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with asthma

When using this product do not use more than directed

Stop use and ask a doctor if

- your child gets nervous, dizzy or sleepless
- symptoms do not get better within 7 days or occur with fever
- cough lasts more than 7 days, comes back, or occurs with fever, rash, or persistent headache. These could be signs of a serious condition.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- mL = milliliter

Age	Dose
children 6 years to under 12 years	10 mL every 4 hours
children 4 years to under 6 years	5 mL every 4 hours
children under 4 years	do not use

Other Information

- each 5 mL contains: sodium 3 mg
- do not use if printed neckband is broken or missing
- store at 20-25°C (68-77°F)
- do not refrigerate
- dosing cup provided

Drug Facts (continued)**Inactive Ingredients**

anhydrous citric acid, edetate disodium, FD&C red #40, avor, glycerin, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol solution, sucralose, xanthan gum

Questions or comments? 1-888-287-1915



Satisfaction guaranteed —
 Or we'll replace it or give you
 your money back. For questions
 or comments or to report an
 undesired reaction or side effect,
 please call 1-888-287-1915.

DISTRIBUTED BY: Wal-Mart Stores,
 Bentonville, AR 72716

**This product is not manufactured or distributed
 by Reckitt Benckiser LLC, owner of the registered
 trademark Mucinex.



PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org



equate™

childrens
MULTI-SYMPOM
COLD
Daytime

AGES 4 TO 12 YEARS

Very Berry
 Flavor Liquid



VALUE PACK

equate™

childrens
MULTI-SYMPOM
COLD
Daytime

Dextromethorphan HBr 5 mg,
 Cough Suppressant
 Guaifenesin 100 mg,
 Expectorant
 Phenylephrine HCl 2.5 mg,
 Nasal Decongestant

Alcohol-Free

AGES 4 TO 12 YEARS

Relieves:

- Stuffy Nose
- Cough
- Chest Congestion
- Breaks Up Mucus

Very Berry
 Flavor Liquid
 Dosing Cup Included

6.8 FL OZ (201mL)



VALUE PACK

equate™

childrens
MULTI-SYMPOM
COLD
Daytime

Dextromethorphan HBr 5 mg,
 Cough Suppressant
 Guaifenesin 100 mg,
 Expectorant
 Phenylephrine HCl 2.5 mg,
 Nasal Decongestant

Alcohol-Free

AGES 4 TO 12 YEARS

Relieves:


- Stuffy Nose
- Cough
- Chest Congestion
- Breaks Up Mucus

Very Berry
 Flavor Liquid
 Dosing Cup Included

6.8 FL OZ (201mL)



839D7 2E C1



NDC 45035-476-40

equate™

Compare to Vicks® DayQuil® Severe active ingredients*

DAYTIME **NON-DROWSY** **SEVERE**

Cold & Flu

Pain Reliever, Fever Reducer, Cough Suppressant, Expectorant, Nasal Decongestant

MAXIMUM STRENGTH

Daytime Relief

- **Acetaminophen** – Headache, fever, sore throat, minor aches & pains
- **Dextromethorphan HBr** – Cough
- **Guaifenesin** – Chest congestion
- **Phenylephrine HCl** – Nasal/sinus congestion & sinus pressure

Alcohol Free

12 FL OZ (355mL)

: 60340 2E F4

PARENTS: Learn about teen medicine abuse
www.StopMedicineAbuse.org

DO NOT USE IF PRINTED NECKBAND IS BROKEN OR MISSING

164152

6 81131 12132 3

Drug Facts

Active ingredients (in each 15 mL)	Purpose
Acetaminophen 325 mg.....	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg.....	Cough suppressant
Guaifenesin 200 mg.....	Expectorant
Phenylephrine HCl 5 mg.....	Nasal decongestant

Uses: temporarily relieves common cold/flu symptoms: ■ sinus congestion and pressure ■ nasal congestion ■ minor aches and pains ■ headache ■ sore throat ■ fever ■ cough due to minor throat and bronchial irritation ■ reduces swelling of nasal passages ■ promotes nasal and/or sinus drainage ■ temporarily restores freer breathing through the nose ■ helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if ■ adult takes more than 4,000 mg of acetaminophen in 24 hours ■ child takes more than 5 doses in 24 hours ■ taken with other drugs containing acetaminophen ■ adult has 3 or more alcoholic drinks every day while using this product
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash
If a skin reaction occurs, stop use and seek medical help right away.

PEEL BACK AT CORNER FOR MORE INFORMATION

: 60340 2E B1

Drug Facts (continued)

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
■ if you have ever had an allergic reaction to this product or any of its ingredients

ESIVE AREA
NO VARNISH • NO TYPE

Drug Facts (continued)



When using this product do not use more than directed

Stop use and ask a doctor if ■ you get nervous, dizzy or sleepless ■ pain, nasal congestion, or cough gets worse or lasts more than 5 days (children) or 7 days (adults) ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions ■ take only as directed – see Overdose warning ■ only use the dose cup provided ■ do not exceed 4 doses per 24 hrs adults & children | 30 mL every 4 hrs

ADHESIVE AREA
NO VARNISH • NO TYPE



NDC 49355-725-40

equate™

Compare to
Vicks® DayQuil®
Cold & Flu active
ingredients*

DAYTIME

NON-DROWSY
ALCOHOL FREE

Cold & Flu

Pain Reliever, Fever Reducer, Cough Suppressant, Nasal Decongestant

Multi-Symptom Relief

- **Acetaminophen** - Aches, fever
- **Dextromethorphan HBr** - Cough
- **Phenylephrine HCl** - Nasal congestion

Antihistamine-Free

12 FL OZ (355mL)

: 65640 2E F5

DO NOT USE IF PRINTED NECKBAND IS BROKEN OR MISSING

164152

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

Drug Facts

Active ingredients (in each 15 mL)	Purpose
Acetaminophen 325 mg.....	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg.....	Cough suppressant
Phenylephrine HCl 5 mg.....	Nasal decongestant

Uses ■ temporarily relieves common cold/flu symptoms:
 ■ cough due to minor throat and bronchial irritation
 ■ nasal congestion ■ minor aches and pains
 ■ sore throat ■ headache ■ fever

Warnings
Liver warnings: This product contains acetaminophen. Severe liver damage may occur if ■ adult takes more than 4,000 mg of acetaminophen in 24 hours ■ child takes more than 5 doses in 24 hours ■ taken with other drugs containing acetaminophen ■ adult has 3 or more alcoholic drinks every day while using this product
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash
 If a skin reaction occurs, stop use and seek medical help right away.

PEEL BACK AT CORNER FOR MORE INFORMATION

: 65640 2E B1

Drug Facts (continued)

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
 ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
 ■ if you have ever had an allergic reaction to this product or any of its ingredients

ADHESIVE AREA
NO VARNISH • NO TYPE

Drug Facts (continued)

Stop use and ask a doctor if ■ you get nervous, dizzy or sleepless ■ pain, nasal congestion or cough gets worse or lasts more than 5 days (children) or 7 days (adults) ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.



If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions ■ take only as directed - see Overdose warning ■ only use the dose cup provided ■ do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 6 hrs
children 6 to under 12 yrs	15 mL every 4 hrs

ADHESIVE AREA
NO VARNISH • NO TYPE





Nighttime Cold & Flu

Drug Facts

Active ingredients (in each 30 mL)	Purpose
Acetaminophen 650 mg.....	Pain reliever/fever reducer
Dextromethorphan HBr 30 mg.....	Cough suppressant
Doxylamine succinate 12.5 mg.....	Antihistamine

Uses

temporarily relieves common cold/flu symptoms:

- cough due to minor throat and bronchial irritation
- sore throat
- headache
- minor aches and pains
- fever
- runny nose and sneezing

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough as occurs with smoking, asthma, or emphysema
- a sodium-restricted diet

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product

- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed - see Overdose warning
 - only use the dose cup provided
 - do not exceed 4 doses per 24 hrs
- | | |
|---------------------------------|-------------------|
| adults & children 12 yrs & over | 30 mL every 6 hrs |
| children 4 to under 12 yrs | ask a doctor |
| children under 4 yrs | do not use |

Other information

- each 30 mL contains: sodium 39 mg
- store at 20-25°C (68-77°F)

Inactive ingredients alcohol, anhydrous citric acid, FD&C blue no. 1, FD&C red no. 40, flavor, high fructose corn syrup, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium citrate

Questions or comments? 1-888-287-1915

Daytime Cold & Flu

Drug Facts

Active ingredients (in each 15 mL)	Purpose
Acetaminophen 325 mg.....	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg.....	Cough suppressant
Phenylephrine HCl 5 mg.....	Nasal decongestant

Uses

temporarily relieves common cold/flu symptoms:

- nasal congestion
- cough due to minor throat and bronchial irritation
- sore throat
- headache
- minor aches and pains
- fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if:

- adult takes more than 4,000 mg of acetaminophen in 24 hours
- child takes more than 5 doses in 24 hours
- with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not use more than directed

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- pain, nasal congestion or cough gets worse or lasts more than 5 days (children) or 7 days (adults)
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed - see Overdose warning
 - only use the dose cup provided
 - do not exceed 4 doses per 24 hrs
- | | |
|---------------------------------|-------------------|
| adults & children 12 yrs & over | 30 mL every 4 hrs |
| children 6 to under 12 yrs | 15 mL every 4 hrs |
| children 4 to under 6 yrs | ask a doctor |
| children under 4 yrs | do not use |

Other information

- each 15 mL contains: sodium 7 mg
- store at 20-25°C (68-77°F)

Inactive ingredients butylated hydroxyanisole, edetate disodium, FD&C yellow no. 6, flavor, glycerin, menthol, monobasic sodium phosphate, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sucrose, xanthan gum

Questions or comments? 1-888-287-1915

equate™



NON-DROWSY
ALCOHOL FREE

Cold & Flu

Pain Reliever, Fever Reducer, Cough Suppressant, Nasal Decongestant

Multi-Symptom Relief

